

The growth of matter perturbations in $f(R)$ models

R. Gannouji*, B. Moraes[†] and D. Polarski[‡]

Lab. de Physique Théorique et Astroparticules, CNRS
Université Montpellier II, France

March 1, 2009

Abstract

We consider the linear growth of matter perturbations on low redshifts in some $f(R)$ dark energy (DE) models. We discuss the definition of dark energy (DE) in these models and show the differences with scalar-tensor DE models. For the $f(R)$ model recently proposed by Starobinsky we show that the growth parameter $\gamma_0 \equiv \gamma(z=0)$ takes the value $\gamma_0 \simeq 0.4$ for $\Omega_{m,0} = 0.32$ and $\gamma_0 \simeq 0.43$ for $\Omega_{m,0} = 0.23$, allowing for a clear distinction from Λ CDM. Though a scale-dependence appears in the growth of perturbations on higher redshifts, we find no dispersion for $\gamma(z)$ on low redshifts up to $z \sim 0.3$, $\gamma(z)$ is also quasi-linear in this interval. At redshift $z = 0.5$, the dispersion is still small with $\Delta\gamma \simeq 0.01$. As for some scalar-tensor models, we find here too a large value for $\gamma'_0 \equiv \frac{d\gamma}{dz}(z=0)$, $\gamma'_0 \simeq -0.25$ for $\Omega_{m,0} = 0.32$ and $\gamma'_0 \simeq -0.18$ for $\Omega_{m,0} = 0.23$. These values are largely outside the range found for DE models in General Relativity (GR). This clear signature provides a powerful constraint on these models.

PACS Numbers: 04.62.+v, 98.80.Cq

*email:gannouji@lpta.univ-montp2.fr

[†]email:moraes@lpta.univ-montp2.fr

[‡]email:polarski@lpta.univ-montp2.fr

1 Introduction

The present stage of accelerated expansion of the universe [1] is a major challenge for cosmology. There are basically two main roads one can take: either to introduce a new (approximately) smooth component or to modify the laws of gravity on cosmic scales. In the first class of models, the new component dubbed dark energy (DE) must have a sufficiently negative pressure in order to induce a late-time accelerated expansion. One typically adds a new isotropic perfect fluid with negative pressure able to accelerate the expansion. In the second class of models one is trying to explain the accelerated expansion by modifying gravity, after all the universe expansion is a large-scale gravitational effect. It is far from clear at the present time which class of DE models will finally prevail and one must study carefully all possibilities. While all models are called DE models [2], the models of the second class are called more specifically modified gravity DE models.

It is clear that in both classes we have many models able to reproduce a late-time accelerated expansion in agreement with the present data probing the background expansion like luminosity distances from SNIa. Probably this will remain the case even with more accurate data. However the evolution of matter perturbations will be affected differently depending on the class of models we are dealing with. This can be used for a consistency check in order to find out whether a DE model is inside General Relativity (GR) or not [3]. Therefore the evolution of matter perturbations seems to be a powerful tool to discriminate between models inside and outside GR [4]. In particular, the growth of matter perturbations on small redshifts seems a promising tool and it has been the subject of intensive investigations recently [5].

A particular class of modified gravity DE models are $f(R)$ models where R is replaced by some function $f(R)$ in the gravitational Lagrangian density [6] (for a recent review, see [7]). The idea of producing an accelerated stage in such models was first successfully implemented in Starobinsky's inflationary model [8]. In some sense it was rather natural to try explaining the late-time accelerated expansion by the same kind of mechanism, a modification of gravity but now at much lower energies. Some problems arising in these $f(R)$ DE models were quickly pointed out, related to solar-system constraints [9] and instabilities [10]. It was then found that a very serious problem arises where it was not expected: many of these models are unable to produce the late-time acceleration together with a viable cosmic expansion history [11]. It was shown that many popular $f(R)$ models, like those containing a R^{-1} term [12], are unable to account for a viable expansion history because of the absence of a standard matter-dominated stage $a \propto t^{\frac{2}{3}}$ which is replaced instead by the behaviour $a \propto t^{\frac{1}{2}}$ [13].

However some interesting cosmological models remain viable like the model recently suggested by Starobinsky [14], (for another interesting viable model see also [15]). It is interesting in the first place because of its non-trivial ability to allow for a standard matter-dominated stage. This goes together with the appearance of large oscillations in the past and the overproduction of massive scalar particles in the very early universe which, as noted already in [14], can be a serious problem of all viable $f(R)$ DE models. Another interesting property is the scale dependence of the

growth of matter perturbations deep inside the Hubble radius [16],[17]. This scale dependence was actually used in [14] in order to constrain one of the parameters of the model. Of course this model also satisfies the local gravity constraints for certain parameter values. In the end a window in parameter space remains where the model is in principle physically acceptable.

In this letter we want to assess the viability of this model with respect to the growth of matter perturbations (see e.g. [18] for other possible constraints and approaches). Some viable $f(R)$ DE models have their free parameters constrained in such a way that they cannot be distinguished observationally from Λ CDM. The situation is different in this case. Our results show that the growth of matter perturbations on low redshifts $z \lesssim 1$ provides a discriminating signature of these models, able to clearly differentiate it from Λ CDM and also from any other DE model inside GR. These results confirm that the growth of matter perturbations can be used efficiently to track the nature of DE models.

2 $f(R)$ modified gravity cosmology

We consider now $f(R)$ models and discuss some general properties in connection with their observational viability. Many $f(R)$ models have the surprising property that they cannot account simultaneously for a viable cosmic expansion history and a late-time accelerated expansion. This severely constraints viable $f(R)$ modified gravity DE models. We consider a universe where gravity and the content of the universe are described by the following action

$$\mathcal{S} = \int d^4x \sqrt{-g} \left[\frac{1}{16\pi G_*} f(R) + \mathcal{L}_m \right]. \quad (1)$$

For the time being, G_* is a bare gravitational constant, its connection with observations depends on the theory under consideration. We concentrate on spatially flat Friedman-Lemaître-Robertson-Walker (FLRW) universes with a time-dependent scale factor $a(t)$ and a metric

$$ds^2 = -dt^2 + a^2(t) d\mathbf{x}^2. \quad (2)$$

For this metric the Ricci scalar R is given by

$$R = 6 \left(2H^2 + \dot{H} \right), \quad (3)$$

where $H \equiv \frac{\dot{a}}{a}$ is the Hubble rate while a dot stands for a derivative with respect to the cosmic time t . DE models have $\ddot{a} > 0$ on low redshifts. The following equations are obtained

$$3FH^2 = 8\pi G_* (\rho_m + \rho_{rad}) + \frac{1}{2}(FR - f) - 3H\dot{F}, \quad (4)$$

$$-2F\dot{H} = 8\pi G_* \left(\rho_m + \frac{4}{3}\rho_{rad} \right) + \ddot{F} - H\dot{F}, \quad (5)$$

where

$$F \equiv \frac{df}{dR} . \quad (6)$$

In standard Einstein gravity ($f = R$) one has $F = 1$. The densities ρ_m and ρ_{rad} satisfy the usual conservation equations $\dot{\rho}_i = -3H(1 + w_i)\rho_i$ with $w_m = 0$ and $w_{rad} = \frac{1}{3}$.

We summarize briefly the way in which the dynamics of $f(R)$ models can be analyzed. For a general $f(R)$ model we introduce the following (dimensionless) variables

$$x_1 = -\frac{\dot{F}}{HF} , \quad (7)$$

$$x_2 = -\frac{f}{6FH^2} , \quad (8)$$

$$x_3 = \frac{R}{6H^2} = \frac{\dot{H}}{H^2} + 2 , \quad (9)$$

$$x_4 = \frac{8\pi G_* \rho_{rad}}{3FH^2} . \quad (10)$$

From Eq. (4) we have the algebraic identity

$$\tilde{\Omega}_m = \frac{8\pi G_* \rho_m}{3FH^2} = 1 - x_1 - x_2 - x_3 - x_4 . \quad (11)$$

It is then possible to write down the following autonomous system

$$\frac{dx_1}{dN} = -1 - x_3 - 3x_2 + x_1^2 - x_1x_3 + x_4 , \quad (12)$$

$$\frac{dx_2}{dN} = \frac{x_1x_3}{m} - x_2(2x_3 - 4 - x_1) , \quad (13)$$

$$\frac{dx_3}{dN} = -\frac{x_1x_3}{m} - 2x_3(x_3 - 2) , \quad (14)$$

$$\frac{dx_4}{dN} = -2x_3x_4 + x_1x_4 , \quad (15)$$

where $N \equiv \ln a$. In eqs(13,14), the quantity m corresponds to

$$m \equiv \frac{d \log F}{d \log R} = \frac{RF'}{F} , \quad (16)$$

(a prime stands for derivative with respect to R) hence m is a priori a function of R . However it is easy to see that R can in turn be expressed in function of the variables of our autonomous system. Indeed we have

$$\frac{x_3}{x_2} \equiv r = -\frac{RF}{f} = -\frac{d \log f}{d \log R} . \quad (17)$$

Inverting (17) we can in principle express R as a function of $\frac{x_3}{x_2} \equiv r$ and so m becomes in turn a function of r , $m = m(r)$ which closes our system.

It was found in [13] that when radiation is negligible ($x_4 = 0$) there exists a critical point $x_1 = 0, x_2 = -\frac{1}{2}, x_3 = \frac{1}{2}$ corresponding to an exact matter phase

$$w_{\text{eff}} = -1 - \frac{2\dot{H}}{3H^2} = \frac{1}{3}(1 - 2x_3) = 0 \quad (18)$$

$$\tilde{\Omega}_m = 1. \quad (19)$$

This is the critical point called $P_5(m = 0)$ in [13], it satisfies $r = -1$.

Let us comment about the physical meaning of these conditions. For $r = -1$ or $x_2 + x_3 = 0$, eq.(4) reduces to

$$3FH^2 = 8\pi G_* \rho_m - 3H\dot{F}. \quad (20)$$

Then we clearly have an exact matter phase ($a \sim t^{\frac{2}{3}}$) for $\dot{F} = 0$ or $x_1 = 0$. We note further from eqs.(13,14) that $x_2 + x_3$ remains zero for $x_1 = 0$.

The condition $m = 0$ is more subtle. From eq.(14) a stable matter phase requires

$$x_1 = 3m. \quad (21)$$

It is straightforward to generalize (21) for arbitrary scaling behaviour of the background

$$x_1 = 3m(1 + w_{\text{eff}}). \quad (22)$$

Actually neither x_1 nor m can be exactly zero during a matter phase (which corresponds to a critical point of the system). This would require $F' = 0$, which reduces to General Relativity plus a cosmological constant. The system can only come in the vicinity of P_5 and trajectories with an acceptable matter phase have $x_1 \approx 0$ and $m \approx 0$ with $x_1 \approx 3m$. In addition, from $F' > 0$ both x_1 and m should be positive during the matter phase. Finally, it is easy to check that (21) implies

$$R \propto a^{-3}. \quad (23)$$

The situation is very different in scalar-tensor models where a standard matter phase is possible with $\dot{F} \approx 0$ because the energy density associated with the dilaton ϕ can mimic the behaviour of dustlike matter for a particular type of potential $U(\phi)$ [19]. It is this property which is used in [20]. More explicitly, for the Lagrangian density

$$\mathcal{L} = \frac{1}{16\pi G_*} \left(F(\Phi) R - g^{\mu\nu} \partial_\mu \Phi \partial_\nu \Phi - 2U(\Phi) \right) + L_m(g_{\mu\nu}), \quad (24)$$

we have the Friedmann equations (in flat space)

$$3FH^2 = 8\pi G_* \rho_m + \frac{\dot{\Phi}^2}{2} + U - 3H\dot{F}, \quad (25)$$

$$-2F\dot{H} = 8\pi G_* \rho_m + \ddot{\Phi} + \ddot{F} - H\dot{F}. \quad (26)$$

It is possible to have a solution with $\dot{F} = 0$ and $\frac{\dot{\Phi}^2}{2} + U \propto a^{-3}$ by a suitable choice of the potential U [19]. Because of this property it is possible in the matter phase

to have a nonvanishing DE energy density ρ_{DE} . The definition of ρ_{DE} requires some care. If we want ρ_{DE} to obey the usual energy conservation for perfect fluids, certainly a desirable property, then the choice is not unique. The Friedmann equations (4,5) and also (25,26) can be recast in the form

$$3A \dot{H}^2 = 8\pi G_* (\rho_m + \rho_{DE}(A)) \quad (27)$$

$$-2A \ddot{H} = 8\pi G_* (\rho_m + \rho_{DE}(A) + p_{DE}(A)) \quad (28)$$

where A is some arbitrary constant. We have written $\rho_{DE}(A)$ (and $p_{DE}(A)$) as these quantities depend on the choice of A . For example from (4,5), neglecting radiation, we obtain

$$8\pi G_* \rho_{DE}(A) = \frac{1}{2}(FR - f) - 3H\dot{F} + 3(A - F) \dot{H}^2 \quad (29)$$

$$8\pi G_* (\rho_{DE}(A) + p_{DE}(A)) = \ddot{F} - H\dot{F} - 2(A - F) \ddot{H} \quad (30)$$

Obviously, whatever choice is adopted the corresponding DE component obeys the usual conservation law of an isotropic perfect fluid. For the representation (27), it is natural to introduce the cosmic relative densities

$$\Omega_i \equiv \frac{8\pi G_* \rho_i}{3A\dot{H}^2} \quad (31)$$

A natural choice, especially if we are willing to compare modified gravity theories with General Relativity (GR), is to take

$$\frac{G_*}{A} \simeq G_N \quad (32)$$

to very high accuracy, where G_N is Newton's constant found in textbooks. In GR, it is the bare gravitational constant appearing in the action and it corresponds to the value of the gravitational constant obtained in a Cavendish-type experiment measuring the Newtonian force between two close test masses. As is well-known, this identification no longer holds outside GR. For modified gravity theories it is the quantity G_{eff} , a priori different from G_* , which plays the role of gravitational constant and which is measured in a Cavendish-type experiment and clearly all acceptable models must have the same G_{eff} today to very high precision.

For scalar-tensor models we can make the choice $A = F_0$ [19] as $\frac{G_*}{F_0} \simeq G_{\text{eff},0}$ to very high accuracy because $\omega_{BD,0} > 4 \times 10^4$ from solar system constraints. As we will see later, in our $f(R)$ model a Cavendish-type experiment corresponds to the limit $G_{\text{eff}} = G_*$ (see [14] and eq. (50) below). It is therefore natural to take $A = 1$ in such a model.

As noted in [14] a consequence of the arbitrariness in the choice of A is that it introduces an arbitrariness in the sign of ρ_{DE} as well. In a model where $F_0 > F$ in the past, ρ_{DE} remains positive if we take $A = F_0$. In our $f(R)$ models however, we have $F_0 < F(z > 0)$ in an expanding universe with $F \rightarrow 1$ at high redshifts. Hence

$\rho_{DE}(A)$ defined from (27) can become negative in the past if we take $A = F_0$. This problem is avoided if we define ρ_{DE} from (27,28) putting $A = 1$

$$3H^2 = 8\pi G_* (\rho_m + \rho_{DE}) \quad (33)$$

$$-2\dot{H} = 8\pi G_* (\rho_m + \rho_{DE} + p_{DE}) . \quad (34)$$

The corresponding relative densities become $\Omega_i \equiv \frac{8\pi G_* \rho_i}{3H^2}$ and actually $\Omega_{DE} \rightarrow 0$ in the past. Note that though it is always possible to write the Friedmann equations in the form (33,34) this should not conceal the fact that we are dealing with a genuinely modified gravity theory obeying eqs.(4,5). The difference is absorbed in the microscopic definitions of the DE component. This shows again the need to go beyond the background expansion in order to discriminate modified gravity DE models from GR.

3 Starobinsky's model

It was shown that a viable cosmic expansion is rather problematic in $f(R)$ models. As already mentioned in the Introduction problems related to instabilities [10] and solar-system constraints [9] were pointed out soon after the first attempts to produce $f(R)$ DE models. An even more unexpected problem was found later related to the background expansion as many $f(R)$ models, including popular ones containing an inverse power of R , are unable to reproduce a viable expansion history containing a standard matter-dominated stage [11]. The viability of these models was then systematically studied in [13]. All this shows that the construction of a viable $f(R)$ DE model is something highly non-trivial. Still there are some $f(R)$ DE models left that can account for a viable expansion history and still depart from Λ CDM. One such interesting model was suggested recently by Starobinsky

$$f(R) = R + \lambda R_c \left(\left(1 + \frac{R^2}{R_c^2} \right)^{-n} - 1 \right) , \quad (35)$$

with $\lambda, n > 0$. We see that $f(R) \rightarrow R$ for $R \rightarrow 0$ in contrast to models containing an inverse power of R , in addition in flat space the correction term to Einstein gravity disappears. The quantity R_c has the dimension of R and is a free parameter of the model, it corresponds essentially to the present cosmic value of R . Recently problems related to the high-curvature regime of this model were pointed out [21] but it seems this could be cured by the addition of a term $\frac{R^2}{\bar{m}^2}$ as noted in [14]. However the mass $\bar{m}$ is required to be very high with $\bar{m} \gg 1$ GeV hence this term is negligible in the low curvature regime. In the present work we will show that it is the low curvature limit of this model, through the growth of matter perturbations on small redshifts, that could severely constrain this model. It is convenient to introduce the reduced dimensionless curvature $\tilde{R} \equiv \frac{R}{R_c}$. We have for this model

$$r(R) = \frac{2n\lambda\tilde{R} \left(\tilde{R}^2 + 1 \right)^{-n-1} - 1}{1 + \lambda\tilde{R}^{-1} \left(\left(\tilde{R}^2 + 1 \right)^{-n} - 1 \right)} \quad (36)$$

This is hardly invertible but fortunately the situation improves for $\tilde{R} \gg 1$. In this regime we have

$$f(R) \simeq R - \lambda R_c \equiv R - 2\Lambda(\infty) \quad (37)$$

$$F \simeq 1 \quad (38)$$

$$r \simeq -1 - \lambda \tilde{R}^{-1} \quad (39)$$

$$m(r) \simeq \frac{2n(2n+1)}{\lambda^{2n}} (-r-1)^{2n+1} \simeq 2n(2n+1) \frac{\lambda}{\tilde{R}^{2n+1}} \approx 0, \quad m > 0 \quad (40)$$

$$M^2 \simeq \frac{1}{3F'} = \frac{R}{3mF} \gg R. \quad (41)$$

It is clear that when $f(R)$ satisfies (37), $x_2 + x_3$ comes close to zero as $FR - f \simeq 2\Lambda(\infty) \ll 6H^2$. We see that M^2 , the scalaron mass introduced in [8], becomes very large in the past. As noted in [14], one should find a mechanism to avoid M^2 becoming too large in the early universe. We get also strong oscillations δR in the past and we illustrate this with the oscillations induced in the behaviour of w_{eff} on high redshifts already during the matter phase as displayed on Figure 1a. These oscillations could lead to overproduction of scalarons in the very early universe [14]. While the cosmic background today corresponds to $\tilde{R} \sim 1$, the regime $\tilde{R} \gg 1$, eqs.(37)-(41), is quickly reached as we go back in time. In this regime $r \approx -1$ and $m \approx 0$ so that a viable matter phase is possible followed by a late-time accelerated stage. We have also in this regime $F \approx 1$ to very high accuracy. One finds that in this model accelerated expansion starts at $z_a \simeq (0.5 - 1)$ depending on the value of $\Omega_{m,0}$, in agreement with observations [22].

In order to be viable, this theory should not be in conflict with gravity constraints. On scales satisfying today $k \ll 2\pi M$, one will not feel the presence of a “fifth force” caused by the scalar part of the gravitational interaction. Hence provided the scalaron mass M corresponding to some experiment is large enough, the “fifth force” will only be felt on scales that are much smaller than those probed by this experiment. In particular one will not see it in a Cavendish-type experiment on scales larger than $5 \times 10^{-2} \text{cm}$ provided $n \geq 1$, and this range can be increased to even lower scales for larger n [14]. We refer the interested reader to the reference [14] for a thorough discussion of the properties of this model and of its potential problems.

The evolution and growth of matter perturbations can also be seen as an experiment probing gravity. In this case the scalaron mass can be much smaller so that deviations from GR and the appearance of a “fifth force” can be felt on larger scales and even on cosmic scales. We will consider this in more detail in the next section.

4 Linear growth of matter perturbations

In a way analogous to scalar-tensor DE models, the equation governing the growth of matter perturbations on subhorizon scales is of the form [23]

$$\ddot{\delta}_m + 2H\dot{\delta}_m - 4\pi G_{\text{eff}} \rho_m \delta_m = 0. \quad (42)$$




Figure 1: a) On the left panel the effective (total) equation of state parameter w_{eff} is shown for $n = 2$. We see that large oscillations appear in the past, already in the matter-dominated stage at rather low redshifts. b) On the right panel, the behaviour of the quantity $\frac{a\tilde{H}}{a_0 H_0}$ is displayed. It is seen that some subhorizon scales satisfying $k \ll a\tilde{H}$ in the past will gradually shift to the regime $k \gg a\tilde{H}$ as the universe expands. In this asymptotic regime, the scalaron corresponding to the cosmic background curvature is nearly massless and a fifth-force is felt on these cosmic scales. Note that today $\tilde{H}_0 \sim 7 H_0$.

The expression for G_{eff} in $f(R)$ models is given below, see eq.(42). Many DE models outside GR have a modified equation of this type describing the growth of matter perturbations. For the models where it is valid, this equation includes all perturbations, the assumption is that we are in a regime deep inside the Hubble radius where the leading terms result in (42). One can also have in principle more elaborate DE models yielding a different equation (see e.g. [24]). It is convenient to rewrite the corresponding equation satisfied by the quantity $f = \frac{d \ln \delta}{d \ln a}$ [25]

$$\frac{df}{dN} + f^2 + \frac{1}{2} \left(1 - \frac{d \ln \Omega_m}{dN} \right) f = \frac{3}{2} \frac{G_{eff}}{G_*} \Omega_m, \quad (43)$$

where Ω_m is defined from (31) with $A = 1$, and [16]

$$G_{eff} = \frac{G_*}{F} \frac{1 + 4 \frac{k^2 F'}{a^2 F}}{1 + 3 \frac{k^2 F'}{a^2 F}}. \quad (44)$$

The r.h.s. of (43) is given by the expression $4\pi G_{eff} \frac{\rho_m}{H^2}$ which obviously does not depend on (the constant) A so that the evolution of f does not depend on A either. If we write this r.h.s. using the definition (29) $\Omega_m = \frac{8\pi G_* \rho_m}{3AH^2}$ with $A \neq 1$, we will have $4\pi G_{eff} \frac{\rho_m}{H^2} = \frac{3}{2} A \frac{G_{eff}}{G_*} \Omega_m$ and we have in particular $\frac{3}{2} A \frac{G_{eff}}{G_*} \Omega_m = \frac{3}{2} \frac{1 + 4 \frac{k^2 F'}{a^2 F}}{1 + 3 \frac{k^2 F'}{a^2 F}} \tilde{\Omega}_m$, with $\tilde{\Omega}_m$ defined in eq.(11). When gravity is described by GR eq.(43) reduces to eq.(B7) given in [26].

¹The reader should not confuse this quantity with the quantity $f(R)$, we believe such confusion will be easily avoided from the context where they appear.

A crucial point in this model is that G_{eff} is scale dependent

$$G_{\text{eff}} = G_{\text{eff}}(z, k) . \quad (45)$$

In other words, the driving force in eq.(43) introduces a scale dependence in the growth of perturbations. The spatial variation on cosmic scales of G_{eff} increases the growth of matter perturbations as is seen very clearly from Figure 2a. As this increase is rather tightly constrained by observations, so will the model parameter n on which this increase depends [14]. We will return in more details to this important point below.

We can rewrite G_{eff} in the suggestive way

$$G_{\text{eff}} = \frac{G_*}{F} \frac{1 + 4 \frac{k^2}{a^2 \tilde{H}^2}}{1 + 3 \frac{k^2}{a^2 \tilde{H}^2}} \quad (46)$$

$$= \frac{G_*}{F} \left(1 + \frac{\frac{k^2}{a^2 \tilde{H}^2}}{1 + 3 \frac{k^2}{a^2 \tilde{H}^2}} \right) , \quad (47)$$

where

$$\tilde{H}^2 = \frac{F}{F'} \equiv 3FM^2 = \frac{R}{m} . \quad (48)$$

In equation (43), G_{eff} depends on the cosmic curvature R taken from (3). In the limit $\tilde{R} \gg 1$ satisfied by the cosmic curvature R at high redshifts, we have

$$G_{\text{eff}} = G_* \left(1 + \frac{1}{3} \frac{\frac{k^2}{a^2 M^2}}{1 + \frac{k^2}{a^2 M^2}} \right) \quad \tilde{R} \gg 1 . \quad (49)$$

Taking the (inverse) Fourier transform of (49) it is straightforward to recover the corresponding gravitational potential per unit mass $V(r)$. Remembering that a potential $\propto \frac{1}{r}$ in real space yields a k^{-2} term in Fourier space, we recognize in (49) the gravitational potential in real space (per unit mass)

$$V(r) = -\frac{G_*}{r} \left(1 + \frac{1}{3} e^{-Mr} \right) . \quad (50)$$

The quantity $\tilde{H}$ (and $M(R)$) can become small enough with the universe expansion, see Figure 1b, so that some cosmic subhorizon scales can feel a significant fifth-force. As the universe expands, $\tilde{H}$ is rapidly decreasing so that these deviations are felt in Poisson's equation on ever increasing scales. While in the past only scales very much smaller than those corresponding to cosmic scales today could feel deviations from GR, today $\tilde{H}_0 \sim H_0$ and deviations can be felt on essentially all subhorizon scales.

We can distinguish two basic asymptotic regimes

$$G_{\text{eff}} = \frac{4}{3} \frac{G_*}{F} \quad k \gg a\tilde{H} , \quad (51)$$

$$= \frac{G_*}{F} \quad k \ll a\tilde{H} . \quad (52)$$

The quantity $a\tilde{H}$ is rapidly increasing in the past. Hence some cosmic subhorizon scales ($k \gg aH$) will satisfy $k \ll a\tilde{H}$ in earlier times (large redshifts) and switch into the regime $k \gg a\tilde{H}$ on lower redshifts as the universe expands, see Figure 1b. For cosmic scales that are in this regime (51), the scalaron mass appears negligible and a fifth force does appear which results in the factor $\frac{4}{3}$.

An accurate evolution of f or δ_m for arbitrary r.h.s. of eq.(43) requires numerical calculations. But it is possible to find analytically the solutions in the two asymptotic regimes (51,52) during the matter stage when $\Omega_m \approx 1$ and $F \approx 1$. This regime is still valid until low redshifts. When $C \equiv \frac{G_{\text{eff}}}{G_*} \Omega_m = \text{constant}$ we get constant growing mode solutions $f = p$ with (see e.g.[20])

$$p = \frac{1}{4} \left(-1 + \sqrt{1 + 24C} \right) . \quad (53)$$

A similar result can also be obtained in the framework of chameleon models (see e.g. [27]) and inside GR (see e.g. [28]). We have $C \approx 1$ for $k \ll a\tilde{H}$ and $C \approx \frac{4}{3}$ for $k \gg a\tilde{H}$, so that we obtain [14]

$$\delta_m \propto a \quad k \ll a\tilde{H} \quad (54)$$

$$\delta_m \propto a^{\frac{\sqrt{33}-1}{4}} \quad k \gg a\tilde{H} . \quad (55)$$

For subhorizon scales that go from the first into the second regime, this can yield a scale-dependent increase (see Figure 2a) in the growth of matter perturbations. Note that this increase takes place on small cosmic scales. This increase in the growth of matter perturbations will induce a change of shape of the matter power spectrum $P(k)$ inferred from galaxy surveys with a subsequent change of its spectral index n_s^{gal} , $P(k) \propto k^{n_s^{\text{gal}}}$. This can then be compared with the spectral index n_s^{CMB} derived from the CMB anisotropy data resulting in a possible discrepancy between both spectral indices. No significant discrepancy between the two values is allowed by present observations and we have the conservative bound (see e.g. [29])

$$n_s^{\text{gal}} - n_s^{\text{CMB}} < 0.05 \quad (56)$$

So this difference is rather tightly constrained and so is the model parameter n on which it depends. This results in the constraint $n \geq 2$. Because increasing the model parameter n will decrease this discrepancy according to the analytical estimate [14]

$$n_s^{\text{gal}} - n_s^{\text{CMB}} = \frac{\sqrt{33} - 5}{2(3n + 2)} , \quad (57)$$

which is accurate for a wide range of n values (only for large values one should resort to a more accurate numerical estimate), (56,57) result in the constraint $n \geq 2$.

Let us emphasize again at this point the differences with scalar-tensor DE models. In scalar-tensor DE models G_{eff} does not depend on $\mathbf{k}$ (or on $\mathbf{r}$ in real space), at a given time it is the same G_{eff} that enters the equation for the growth of perturbations and the gravitational constant measured in a Cavendish-type experiment. These

DE models have further a negligible dilaton mass but they can comply with solar-system constraints if they satisfy $\omega_{BD,0} > 4 \times 10^4$, where $\omega_{BD,0}$ is the value today of the Brans-Dicke parameter, which yields $\gamma_{PN} \approx 1$ [23],[30]. Indeed for scalar-tensor models one has for the Post-Newtonian parameter today $\gamma_{PN} = \frac{\omega_{BD,0}+1}{\omega_{BD,0}+2}$ and solar-system constraints point at $\gamma_{PN} \approx 1$ close to its GR value $\gamma_{PN} = 1$. As $f(R)$ models correspond to scalar-tensor models with $\omega_{BD} = 0$ they have $\gamma_{PN} = \frac{1}{2}$ which requires a high scalaron mass to comply with solar-system constraints.

In asymptotically stable scalar-tensor DE models the matter perturbations grow slower than a in the matter stage and this growth is scale-independent.

The growth of matter perturbations could provide an efficient way to discriminate between modified gravity DE models and DE models inside GR. One can characterize the growth of matter perturbations on small redshifts using the quantity $\gamma(z)$

$$f = \Omega_m(z)^{\gamma(z)} . \quad (58)$$

In the pioneering papers on this approach γ was taken constant [31]. But it is important to realize that $\gamma(z)$ is a function of z which is generically not constant, and that its variation can contain crucial information about the underlying model. As was shown earlier, for a wide class of models inside GR one has $|\gamma'_0 \equiv \frac{d\gamma}{dz}(z=0)| \lesssim 0.02$ so that $\gamma(z)$ is approximately constant [25]. In Λ CDM we have $\gamma_0 \approx 0.55$ (with a slight dependence on $\Omega_{m,0}$) and $\gamma'_0 \simeq -0.015$. However, γ'_0 can be significantly larger in models outside GR [20]. From the growth of matter perturbations we can calculate $\gamma(z)$ and $\frac{d\gamma}{dz}$ and in particular their value at $z=0$. As we can see from Figure 2a, we find that f is scale-independent on low redshifts $z \lesssim 0.5$. This means that the growth of matter perturbations is the same on all relevant scales in this redshift interval. This is easily understood as $\tilde{H}_0 \sim H_0$ so that all subhorizon scales, and certainly the relevant scales that are deep inside the Hubble radius today, satisfy $k \gg \tilde{H}_0$ (we take $a_0 = 1$). Some scale dependence can appear in the growth of matter perturbations *today* for scales that are so large that the original equation (43) (or (42)) no longer provides an accurate approximation. Therefore the quantity $\gamma(z)$ is essentially scale independent too on low redshifts $z \leq 0.3$ in our model. On higher redshifts some restricted dispersion appears which could be another signature of this model, for example at $z = 0.5$ we can have a difference $\Delta\gamma \sim 0.04$ between various scales, see Figure 2b. On even higher redshifts the growth is of course scale-dependent and we can see from Figures 2a that the relative increase is damped with increasing n in accordance with earlier analytical estimates. As mentioned earlier, this increase is constrained by the observations and cannot be large. We find further that in this $f(R)$ model $\gamma_0 \equiv \gamma(z=0) \approx 0.41$ which is much lower than in Λ CDM where $\gamma_0 \approx 0.55$ (and $\gamma'_0 \approx -0.015$). This is an interesting property which clearly allows to discriminate this model from Λ CDM. It is also significantly lower than the value found in some scalar-tensor DE models. This value seems essentially independent of the model parameter n . So a measurement of γ_0 could allow to discriminate this model from Λ CDM, but also possibly from other modified gravity DE models. As γ_0 is scale independent this is a clear signature of this $f(R)$ model.

These conclusions remain the same if we let $\Omega_{m,0}$ vary in the viable range of $\Omega_{m,0}$ values. As we can see on Figures 3, there is some variation of γ_0 and γ'_0 in function




Figure 2: a) On the left, the growth factor f is shown for the model with $n = 2$ and the dimensionless model parameter λ has the value $\lambda = 0.94$. We see the scale dependent behaviour of f . On low redshifts all the cosmic scales displayed here are in the regime (51), and matter perturbations obey (55) in the matter-dominated stage. A significant fifth-force is felt in the deviation of Poisson's equation for matter perturbations on these cosmic scales. Note that on higher redshifts these scales will eventually be in the opposite regime (52), so that perturbations follow the evolution (54). This is the origin of the bumps seen on the figure. b) On the right, the quantity $\gamma(z)$ is displayed for the same model parameters as on the left panel. We find a low value for the growth parameter γ_0 , substantially lower than in Λ CDM. It is seen that the dispersion of $\gamma(z)$ is very small, while that of γ_0 is negligible. We find also a value for $\gamma'_0 \equiv \frac{d\gamma}{dz}(z=0)$, $\gamma'_0 \sim 0.2$ which is much larger than the value found for DE models inside GR. c) On the bottom figure, the quantity $\frac{\delta_m(z)}{\delta_{m,0}}$ is displayed up to $z = 0.5$. This quantity does not depend explicitly on the background quantity Ω_m . At $z = 0.5$, a difference of about 6% is found with Λ CDM.

$\Omega_{m,0}$	0.322	0.315	0.302	0.289	0.273	0.263	0.254	0.245	0.238	0.227
γ_0	0.396	0.399	0.404	0.409	0.415	0.419	0.422	0.425	0.428	0.432
γ'_0	-0.253	-0.246	-0.234	-0.224	-0.210	-0.202	-0.195	-0.189	-0.183	-0.175
z_a	0.654	0.673	0.711	0.747	0.798	0.830	0.862	0.893	0.922	0.965
$\frac{\delta_m(z=0.5)}{\delta_{m,0}}$	0.731	0.733	0.737	0.742	0.747	0.751	0.754	0.758	0.761	0.765

Table 1: This table summarizes the value of the growth parameters γ_0 , γ'_0 and of z_a , the redshift when accelerated expansion starts. All these values correspond to $n = 2$ and $\lambda = 0.94$.




Figure 3: a) On the left, the parameter γ_0 is displayed in function of the cosmological parameter $\Omega_{m,0}$ for different values of the model parameter n . For given n , we see some variation with the highest value obtained for a low matter density universe. Still all these values remain clearly distinguishable from their values in Λ CDM. b) On the right, the parameter γ'_0 . We note that very large values are obtained for all models.

of $\Omega_{m,0}$. We get $\gamma_0 \simeq 0.4$ for $\Omega_{m,0} = 0.32$ and $\gamma_0 \simeq 0.43$ for $\Omega_{m,0} = 0.23$. As for γ'_0 we have $\gamma'_0 \simeq -0.18$ for $\Omega_{m,0} = 0.23$ and $\gamma'_0 \simeq -0.25$ for $\Omega_{m,0} = 0.32$. So a low matter density universe brings the quantities γ_0 and γ'_0 closer to their values in Λ CDM but they still remain far from them. Such low values for γ_0 seem in trouble with present observations (see e.g. [33]) but an appropriate analysis of the data should relax the assumption $\gamma = \text{constant}$. It is likely that such a low value will remain in tension with observations.

Finally we have here again an example of a modified gravity model where $\frac{d\gamma}{dz}(z=0) \equiv \gamma'_0 \neq 0$ and actually large. We obtain the high value $\gamma'_0 \approx -0.14$ which is largely outside the range $|\gamma'_0| \leq 0.02$ found for DE models inside GR. It is also much higher than the value found for some scalar-tensor DE models. So we have here again a characteristic signature of our model which clearly differentiates it both from Λ CDM and DE models inside GR, but possibly also from other DE models outside GR. Like for γ_0 , the scale independence of γ'_0 makes it a clear signature of the model. We have

checked that γ'_0 obeys the constraint

$$\gamma'_0 = [\ln \Omega_{m,0}^{-1}]^{-1} \left[-\Omega_{m,0}^{\gamma_0} - 3\left(\gamma_0 - \frac{1}{2}\right)w_{\text{eff},0} + \frac{3}{2} \frac{G_{\text{eff}}(R_0)}{G_*} \Omega_{m,0}^{1-\gamma_0} - \frac{1}{2} \right], \quad (59)$$

where we have to include the factor $\frac{G_{\text{eff}}(R_0)}{G_*} \simeq \frac{4}{3F_0} \neq 1$. As for the scalar-tensor models considered earlier, we obtain a nearly linear behaviour on low redshifts $z \leq 0.3$.

In conclusion we find that the growth of matter perturbations on small redshifts provides a powerful constraint on our $f(R)$ model. We find low values for the parameter γ_0 with $\gamma_0 \approx 0.4$, and high values for γ'_0 with $\gamma'_0 \simeq -0.2$. These values are not scale dependent and provide therefore a clear signature. The growth parameters γ_0 and γ'_0 are mainly affected by the cosmological parameter $\Omega_{m,0}$, in particular γ'_0 is less negative for low $\Omega_{m,0}$ values. But in all cases, characteristic values very far from Λ CDM and from all DE models inside GR are found.

There are still very large uncertainties on the quantity $f(z)$ or on $\beta \equiv \frac{f}{b}$ where b is the bias factor (see e.g. [33],[32]). One should further keep in mind that a precise observational determination of both $f(z)$ and Ω_m is needed in order to measure $\gamma(z)$ accurately. Hence to get precise values for the couple γ_0, γ'_0 we need to determine accurately both $f(z)$ and $\Omega_m(z)$ around $z = 0$. If future surveys will constrain γ_0, γ'_0 to be close to their values for Λ CDM then the models we have investigated here will be ruled out. Though a systematic numerical exploration in the model parameter space is very hard to achieve, the model for $n = 2$ shows already a very large deviation from Λ CDM if one considers the growth of matter perturbations while this model meets all other constraints. So either the model will be ruled out for any value of n or the growth of matter perturbations will at least significantly restrict the viable interval in the model parameter n .

We conjecture that many if not all viable $f(R)$ models will have similar observational signatures. A precise determination of the parameters γ_0 and γ'_0 could be decisive in the quest for the true DE model especially if it is an $f(R)$ modified gravity DE model.

References

- [1] S.J. Perlmutter et al., Ap. J. **483** 565 (1997), Nature **391** 51 (1998); A. G. Riess, A. V. Filippenko, P. Challis *et al.*, Astron. J. **116**, 1009 (1998); S. J. Perlmutter, G. Aldering, G. Goldhaber *et al.*, Astroph. J. **517**, 565 (1999); P. Astier, J. Guy, N. Regnault *et al.*, Astron. Astroph. **447**, 31 (2006); Adam G. Riess *et al.* e-Print astro-ph/0611572.
- [2] V. Sahni, A. A. Starobinsky, Int. J. Mod. Phys. D**9**, 373 (2000); T. Padmanabhan, Phys. Rep. **380**, 235 (2003); E. J. Copeland, M. Sami and S. Tsujikawa, hep-th/0603057 (2006); V. Sahni, A. A. Starobinsky, Int. J. Mod. Phys. D**15**, 2105 (2006).
- [3] A. A. Starobinsky, JETP Lett.**68**, 757 (1998).

- [4] S. Bludman, e-Print [astro-ph/0702085](#). D. Polarski, AIP Conf. Proc. 861, 1013 (2006), e-Print [astro-ph/0605532](#); M. Ishak, A. Upadhye, D. N. Spergel, Phys. Rev. D **74**, 043513 (2006).
- [5] D. Huterer, E. Linder, [astro-ph/0608681](#); E. Linder, R. Cahn, [astro-ph/0701317](#); A. F. Heavens, T. D. Kitching, L. Verde, e-Print [astro-ph/0703191](#); C. Di Porto, L. Amendola, [arXiv:0707.2686](#); V. Acquaviva, L. Verde, [arXiv:0709.0082](#); A. Kiakotou, O. Elgaroy, O. Lahav, [arXiv:0709.0253](#); Y. Wang, [arXiv:0710.3885](#); L. Hui, K. Parfree, [arXiv:0712.1162](#); J.B. Dent, S. Dutta, [arXiv:0808.2689](#); H. Zhang, H. Yu, H. Noh, Z.-H. Zhu, Phys. Lett. B**665** (2008); Hao Wei, S. N. Zhang, Phys. Rev. D**78**, 023011 (2008); V. Acquaviva, A. Hajian, D. N. Spergel, S. Das, Phys. Rev. D**78**, 043514 (2008); Yungui Gong, [arXiv:0808.1316](#).
- [6] S. Capozziello, Int. J. Mod. Phys. D **11**, 483 (2002).
- [7] T. P. Sotiriou, V. Faraoni, [arXiv:0805.1726](#).
- [8] A. A. Starobinsky, Phys. Lett. B**91**, 99 (1980).
- [9] T. Chiba, Phys. Lett. B **575**, 1 (2003).
- [10] A. D. Dolgov and M. Kawasaki, Phys. Lett. B **573**, 1 (2003).
- [11] L. Amendola, D. Polarski, S. Tsujikawa, Phys. Rev. Lett. **98**, 131302 (2007).
- [12] S. M. Carroll, V. Duvvuri, M. Trodden, M. S. Turner, Phys. Rev. D**70**, 043528 (2004).
- [13] L. Amendola, R. Gannouji, D. Polarski, S. Tsujikawa, Phys. Rev. D**75**, 083504 (2007); L. Amendola, D. Polarski, S. Tsujikawa, Int. J. Mod. Phys. D**16**, 1555 (2007).
- [14] A. A. Starobinsky, JETP Lett.**86**, 157 (2007).
- [15] W. Hu, I. Sawicki, Phys. Rev. D**76**, 064004 (2007).
- [16] S. Tsujikawa, Phys. Rev. D**76**, 023514 (2007).
- [17] E. Bertschinger, P. Zukin, Phys. Rev. D**78**, 024015 (2008).
- [18] A. Dev, D. Jain, S. Jhingan, S. Nojiri, M. Sami, I. Thongkool, [arXiv:0807.3445](#); L. Pogosian, A. Silvestri, Phys. Rev. D **77**, 023503 (2008)
- [19] R. Gannouji, D. Polarski, A. Ranquet, A. A. Starobinsky, JCAP **0609**, 016 (2006).
- [20] R. Gannouji, D. Polarski, JCAP **0805**, 018 (2008).
- [21] V. A. Frolov Phys. Rev. Lett. **101**, 061103 (2008); T. Kobayashi, Kei-ichi Maeda, Phys. Rev. D**78**, 064019 (2008).

- [22] A. Melchiorri, L. Pagano, S. Pandolfi, Phys. Rev. D**76**, 041301 (2007).
- [23] B. Boisseau, G. Esposito-Farèse, D. Polarski and A.A. Starobinsky, Phys. Rev. Lett. **85**, 2236 (2000).
- [24] M. Kunz, D. Sapone, Phys. Rev. Lett. **98**, 121301 (2007).
- [25] D. Polarski, R. Gannouji, Phys. Lett.B**660**, 439 (2008).
- [26] L. Wang, P. J. Steinhardt, Astrophys. J.**508**, 483 (1998).
- [27] P. Brax, C. van de Bruck, A.-C. Davis, Phys. Rev. D**70**, 123518 (2004).
- [28] J. Lesgourgues, S. Pastor, Phys. Rept. **429**, 307 (2006).
- [29] M. Tegmark, D. Eisenstein, M. Strauss, et al, Phys. Rev. D**74**, 123507 (2006).
- [30] G. Esposito-Farèse and D. Polarski, Phys. Rev. D**63**, 063504 (2001).
- [31] P. J. E. Peebles, Astrophys. J.**284**, 439 (1984); O. Lahav, P. B. Lilje, J. R. Primack, M. J. Rees, MNRAS **251**, 128 (1991).
- [32] S. Nesseris and L. Perivolaropoulos, Phys. Rev. D**77**, 023504 (2008).
- [33] K. Yamamoto, T. Sato, G. Huetsi, [arXiv:0805.4789](#).